



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Intraoperative low-tidal-volume ventilation

Mynbaev, Ospan A ; Biro, Peter ; Stark, Michael

DOI: <https://doi.org/10.1056/NEJMc1311316SA3>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-84946>

Journal Article

Published Version

Originally published at:

Mynbaev, Ospan A; Biro, Peter; Stark, Michael (2013). Intraoperative low-tidal-volume ventilation. New England Journal of Medicine, 369(19):1862.

DOI: <https://doi.org/10.1056/NEJMc1311316SA3>

Joseph E. Schwartz, Ph.D.

Stony Brook University
Stony Brook, NY

Since publication of their article, the authors report no further potential conflict of interest.

1. Dadhania D, Snopkowski C, Ding R, et al. Validation of non-invasive diagnosis of BK virus nephropathy and identification of prognostic biomarkers. *Transplantation* 2010;90:189-97.
2. Steyerberg EW, Vickers AJ, Cook NR, et al. Assessing the

performance of prediction models: a framework for traditional and novel measures. *Epidemiology* 2010;21:128-38.

3. Bustin SA, Benes V, Garson JA, et al. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009;55:611-22.
4. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 2008;3:1101-8.
5. Røge R, Thorsen J, Tørring C, Ozbay A, Møller BK, Carstens J. Commonly used reference genes are actively regulated in vitro stimulated lymphocytes. *Scand J Immunol* 2007;65:202-9.

DOI: 10.1056/NEJMc1310006

Intraoperative Low-Tidal-Volume Ventilation

TO THE EDITOR: We are troubled by the ventilation strategy selected for the control group (or nonprotective-ventilation group) in the study by Futier et al. (Aug. 1 issue).¹ This strategy (nonprotective ventilation with a tidal volume of 10 to 12 ml per kilogram of predicted body weight, with no positive end-expiratory pressure [PEEP] and no recruitment maneuvers) is known to be potentially harmful and is outdated (the authors cite a study from 1963² to define their standard of care). The tidal volumes recommended in contemporary strategies^{3,4} for perioperative ventilation are less than 10 ml per kilogram of predicted body weight, and they are provided with PEEP.

To analyze practice patterns for patients undergoing anesthesia, we undertook a study in which we reviewed data from 230,386 surgical procedures at two institutions. These data show that PEEP with a median of 5 cm of water was used in 60.4% of procedures. Median tidal volumes declined from 9.2 to 7.9 ml per kilogram of predicted body weight between 2005 and 2013.

Prospective studies such as that by Futier et al. are most relevant if they derive their data from recent observational studies that used contemporary standards of care. Moreover, these data can be derived from clinical-decision support databases.⁵ Do the authors have contemporary data to support their choice of ventilation strategies?

Jonathan P. Wanderer, M.D.

Vanderbilt University Medical Center
Nashville, TN
jonathan.wanderer@vanderbilt.edu

James M. Blum, M.D.

University of Michigan
Ann Arbor, MI

Jesse M. Ehrenfeld, M.D., M.P.H.

Vanderbilt University Medical Center
Nashville, TN

No potential conflict of interest relevant to this letter was reported.

1. Futier E, Constantin JM, Paugam-Burtz C, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *N Engl J Med* 2013;369:428-37.
2. Bendixen HH, Hedley-Whyte J, Laver MB. Impaired oxygenation in surgical patients during general anesthesia with controlled ventilation: a concept of atelectasis. *N Engl J Med* 1963;269:991-6.
3. The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342:1301-8.
4. Wolthuis EK, Choi G, Delsing MC, et al. Mechanical ventilation with lower tidal volumes and positive end-expiratory pressure prevents pulmonary inflammation in patients without preexisting lung injury. *Anesthesiology* 2008;108:46-54.
5. Blum JM, Stentz MJ, Maile MD, et al. Automated alerting and recommendations for the management of patients with preexisting hypoxia and potential acute lung injury: a pilot study. *Anesthesiology* 2013;119:295-302.

DOI: 10.1056/NEJMc1311316

TO THE EDITOR: Futier et al. attributed the outcome of fewer postoperative complications in the protective-ventilation group, in which low-tidal-volume ventilation and PEEP were used, mainly to the prevention of atelectasis. We postulate that microaspiration could be another reason for the higher rate of postoperative pneumonia in the nonprotective-ventilation group.

In our recent study,¹ we found that without the provision of PEEP there was downward leakage of fluid across the cuff of the endotracheal tube, even at a recommended cuff pressure of 20 to 30 cm of water. This leakage was eliminated with the use of a PEEP as low as 5 cm of water or with the use of endotracheal tubes with newer cuff designs. In another study² in which high-tidal-volume ventilation was compared with low-tidal-volume ventilation, with a PEEP of 5 cm of water in both groups during major upper abdominal surgery, no significant difference was detected in postoperative lung function or clinical

cal outcome. We believe the application of PEEP is the critical variable in the study by Futier et al. We also wonder whether the inflammatory condition associated with pneumonia contributed to the anastomotic leak.

Sin-Man Lam, M.D.
Arthur C.W. Lau, M.D.
Kenny K.C. Chan, M.D.

Pamela Youde Nethersole Eastern Hospital
Hong Kong, China
lamsm2@ha.org.hk

No potential conflict of interest relevant to this letter was reported.

1. Lau ACW, Lam SM, Yan WW. Benchtop study of leakages across the Portex, TaperGuard and Microcuff endotracheal tubes under simulated clinical conditions. *Hong Kong Med J* 2013 July 22 (Epub ahead of print).
2. Treschan TA, Kaisers W, Schaefer MS, et al. Ventilation with low tidal volumes during upper abdominal surgery does not improve postoperative lung function. *Br J Anaesth* 2012;109:263-71.

DOI: 10.1056/NEJMc1311316

TO THE EDITOR: Futier et al. compared two approaches to ventilation during laparoscopic and open surgery (laparotomy). Primary outcomes were recorded in 69 of the 85 patients undergoing laparoscopic surgery and in 7 of 315 patients undergoing nonlaparoscopic open surgery. The primary strategies used in open and laparoscopic surgical ventilation support are different, partly because of the insufflation of carbon dioxide that occurs during laparoscopic surgery.¹⁻⁴ To avoid hypercapnia and respiratory acidosis during laparoscopic surgery, it is common to use gradually increasing tidal volumes, ultimately leading to the use of high-volume–high-pressure mechanical ventilation during prolonged laparoscopic surgeries. Tidal volumes can be adjusted by monitoring the end-tidal carbon dioxide concentration and the acid–base status of the blood. To what extent were the ventilator settings changed during laparoscopic surgery to meet its unique needs? Can the lack of such adjustments explain the observed outcomes?

Ospan A. Mynbaev, M.D., Ph.D.
Moscow Institute of Physics and Technology
Moscow, Russia

Peter Biro, M.D.
University Hospital Zurich
Zurich, Switzerland

Michael Stark, M.D.
New European Surgical Academy
Berlin, Germany
mstark@nesacademy.org

No potential conflict of interest relevant to this letter was reported.

1. Neudecker J, Sauerland S, Neugebauer E, et al. The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery. *Surg Endosc* 2002;16:1121-43.
2. Agresta F, Ansaloni L, Baiocchi GL, et al. Laparoscopic approach to acute abdomen from the Consensus Development Conference of the Società Italiana di Chirurgia Endoscopica e nuove tecnologie (SICE), Associazione Chirurghi Ospedalieri Italiani (ACOI), Società Italiana di Chirurgia (SIC), Società Italiana di Chirurgia d'Urgenza e del Trauma (SICUT), Società Italiana di Chirurgia nell'Ospedalità Privata (SICOP), and the European Association for Endoscopic Surgery (EAES). *Surg Endosc* 2012;26:2134-64.
3. Nguyen JH, Tanaka PP. Chapter 15: anesthesia for laparoscopic surgery. In: Wetter PA, Kavic MS, Nezhat C, Winfield H, Kelley WE Jr, Mettler L, eds. *Prevention and management of laparoendoscopic surgical complications (PM4.0)*. 3rd ed. Miami: Society of Laparoendoscopic Surgeons, 2012 (http://laparoscopy.blogs.com/prevention_management_3/2010/10/anesthesia-for-laparoscopic-surgery.html).
4. Russo A, Di Stasio E, Scagliusi A, et al. Positive end-expiratory pressure during laparoscopy: cardiac and respiratory effects. *J Clin Anesth* 2013;25:314-20.

DOI: 10.1056/NEJMc1311316

THE AUTHORS REPLY: In response to Wanderer et al.: we fully concur that there has been a gradual decrease over time in the use of high-tidal-volume ventilation and that the ventilation strategy used in our control group may therefore not fully reflect practice in all hospitals. Nevertheless, lung protection is often incorrectly reduced to lowering tidal volume, with either very low levels of PEEP (<5 cm of water) or no PEEP. Low-tidal-volume ventilation alone is not only ineffective¹ but also may be deleterious. Contemporary data on the use of PEEP in addition to lower tidal volume are scarce. A recent analysis of an electronic database from the University of Colorado suggested that a combination of a tidal volume of more than 8 ml per kilogram of predicted body weight and a PEEP of less than 5 cm of water represented the standard of care in more than 60% of patients.² Data from our observational study indicate that regardless of the PEEP level used, a tidal volume in the range of our protective-ventilation strategy (6 to 8 ml of water per kilogram of predicted body weight) was used in 30.1% of patients.³

We fully agree with the comment by Lam et al. on the pivotal role of PEEP in addition to lower tidal volume in the protective-ventilation strategy. However, the additional contribution of recruitment maneuvers (which were a part of this strategy) should not be underestimated, since PEEP levels in the range of those used in our

experimental conditions are not effective in re-opening nonaerated lung regions.⁴ Further investigation will be required to address the question of whether the inflammatory conditions associated with the development of pneumonia or the use of mechanical ventilation can explain anastomotic leak.

Mynbaev et al. ask about the possible influence of laparoscopic surgery on the occurrence of the primary outcome. Data presented for laparoscopic surgery in Table S1 of the Supplementary Appendix of our article (available at NEJM.org) were erroneously interchanged with the data for nonlaparoscopic surgery. The primary outcome measure should have been shown for 10 of the 85 patients who underwent laparoscopic surgery rather than for 69 of those 85 patients (odds ratio, 0.50; 95% confidence interval, 0.25 to 1.03; $P=0.06$) and for 66 of the 315 patients who underwent nonlaparoscopic surgery. We regret the error; a corrected version of the Supplementary Appendix is available at NEJM.org.

Emmanuel Futier, M.D., Ph.D.

Bruno Pereira, Ph.D.

Centre Hospitalier Universitaire Estaing
Clermont-Ferrand, France

Samir Jaber, M.D., Ph.D.

Centre Hospitalier Universitaire Saint-Eloi
Montpellier, France
s-jaber@chu-montpellier.fr

Since publication of their article, the authors report no further potential conflict of interest.

1. Treschan TA, Kaisers W, Schaefer MS, et al. Ventilation with low tidal volumes during upper abdominal surgery does not improve postoperative lung function. *Br J Anaesth* 2012;109:263-71.
2. Fernandez-Bustamante A, Wood CL, Tran ZV, Moine P. Intraoperative ventilation: incidence and risk factors for receiving large tidal volumes during general anesthesia. *BMC Anesthesiol* 2011;11:22.
3. Jaber S, Coisel Y, Chanques G, et al. A multicentre observational study of intra-operative ventilatory management during general anaesthesia: tidal volumes and relation to body weight. *Anaesthesia* 2012;67:999-1008.
4. Duggan M, Kavanagh BP. Pulmonary atelectasis: a pathogenic perioperative entity. *Anesthesiology* 2005;102:838-54.

DOI: 10.1056/NEJMc1311316

Glucose Levels and Risk of Dementia

TO THE EDITOR: Crane and coworkers (Aug. 8 issue)¹ report on the relationship between glucose levels and the risk of dementia in participants whose mean age at baseline was 76 years. Their analysis is limited by the fact that the established and readily available confounder of renal function had not been accounted for. In the United States, 62.2% of persons 80 years of age or older have an estimated glomerular filtration rate (GFR) of less than 60 ml per minute per 1.73 m² of body-surface area.² Moderate renal impairment has been shown to be associated with an excess risk of incident dementia among persons in good-to-excellent health.³ A community-based cross-sectional study showed that global performance and specific cognitive functions are negatively affected early in chronic kidney disease.⁴ In a community-based study with a follow-up period similar to that in the study of Crane et al., the change in renal function over time was related to the change observed in global cognitive ability, verbal episodic memory, and abstract reasoning.⁵ Aside from the estimated GFR, increased albuminuria is also independently associated with a faster decline in cognitive function. Hence, renal function is an important factor contributing to cognitive impairment and cognitive decline.

Jan T. Kielstein, M.D.

Hannover Medical School
Hannover, Germany
kielstein@yahoo.com

No potential conflict of interest relevant to this letter was reported.

1. Crane PK, Walker R, Hubbard RA, et al. Glucose levels and risk of dementia. *N Engl J Med* 2013;369:540-8.
2. Grams ME, Juraschek SP, Selvin E, et al. Trends in the prevalence of reduced GFR in the United States: a comparison of creatinine- and cystatin C-based estimates. *Am J Kidney Dis* 2013; 62:253-60.
3. Seliger SL, Siscovick DS, Stehman-Breen CO, et al. Moderate renal impairment and risk of dementia among older adults: the Cardiovascular Health Cognition Study. *J Am Soc Nephrol* 2004; 15:1904-11.
4. Elias MF, Elias PK, Seliger SL, Narsipur SS, Dore GA, Robbins MA. Chronic kidney disease, creatinine and cognitive functioning. *Nephrol Dial Transplant* 2009;24:2446-52.
5. Davey A, Elias MF, Robbins MA, Seliger SL, Dore GA. Decline in renal functioning is associated with longitudinal decline in global cognitive functioning, abstract reasoning and verbal memory. *Nephrol Dial Transplant* 2013;28:1810-9.

DOI: 10.1056/NEJMc1311765

THE AUTHORS REPLY: Kielstein raises the possibility that renal function may confound the association that we found between glucose levels and dementia risk. We are aware of the literature that Kielstein references. Indeed, these data pro-